

280 x 100

CAUTION:

No incompatibility at the prescribed doses. WAITING PERIOD for meat and milk: 2 days.

While using product, contact with skin and mucosa must be avoided. After use, wash hands and exposed body parts with soap and water. In case of accidental ingestion, provoke vomit and see a doctor taking the container with you.

**VETERINARY USE
MADE IN URUGUAY**

Batch N°:

Fabrication date:

Expiration date:

Reg. M.G.A.P. N° A-1251

Owner and manufacturer:

LABORATORIOS MICROSULES URUGUAYS S.A.

Ruta 101 km.28 - Cno. al Paso Escobars/n

Pueblo Cap. J. A. Artigas - Canelones

Uruguay - C.P. 91.001

Tel. (00598) 2288 6761*

Fax (00598) 2288 6760

Technical Director: Dr. Pablo R. Bringas

www.laboratoriosmicrosules.com



**CONTENT: 1 Lt.
VETERINARY USE**

Ectosules 15 %



CYPERMETHRIN 15%
EMULSIFYABLE CONCENTRATED
FOR BOVINE AGAINST TICKS AND INSECTICIDE

MICROSULES

FORMULA:

Each 100 ml contains:

Cypermethrin..... 15 g

Excipients q.s.ad.100 ml

MT2333
03/03/2017 - V01

INDICATIONS: External parasiticide

Long-lasting residual action (8 to 14 days) against tick. Repellent action insecticide for a long term period between 3 and 4 weeks.

Mites and ticks: *Boophilus microplus*; *Rhipicephalus sanguineus*; *Psoroptes* spp.; *Sarcoptes scabiei*; *Notoedres cati*.

Flies and Horseflies: *Musca* spp.; *Haematobia irritans* (horn fly); *Callitroga hominivorax*; *Sarcophaga carmaria*; *Lucilia sericata*; *Tabanus* spp.; *Dermatobia hominis*.

Louises: *Damalinea Bovis*; *Damalinea ovis*; *Bovicola subrostratus*; *Haematopinus eurytenuis*; *Haematopinus asini*; *Haematopinus suis*.

Fleas: *Ctenocephalides felis*; *Ctenocephalides canis*.

DOSAGE:

For immersion baths: * Foot of bath: 1 litre Ectosules every 1,000 litres of water * Refill: 1.5 litres of Ectosules every 1,000 litres of water to refill. * Dry reinforcement: 0.5 litres of Ectosules every time the water level in the bath decreases 1,000 litres.

Aspersión: *1 litre of Ectosules every 1,000 litres of water, or 1 unit of 20 ml of Ectosules every 20 lts of water (apply the final emulsion by aspersión using 1 litre per animal).

For pulverisation in premises: *1 litre of Ectosules every 2,00 litres of water or 1 unit of 20 ml of Ectosules every 4 lts of water (proportion: 1 litre of the final emulsion every 10 to 15 m² of surface to treat). For the adequate use of the product, prepare an emulsion in a volume of water equivalent to twice the amount of product to be used.

For strategic treatments in sanitation areas its use is recommended every 21 days.

ROUTE OF ADMINISTRATION:

Used in animals as an immersion bath or aspersión. In premises: by pulverisation.

CAMBIOS 03/03/2017
Corregir Version